

Hirzebruch genera of manifolds equipped with a Hamiltonian circle action

K. E. Feldman*

In the paper we describe obstructions for the existence of symplectic and Hamiltonian symplectic circle actions on closed compact manifolds in terms of Hirzebruch genera and relations between differential and homotopic invariants of such manifolds. All manifolds considered are assumed to be closed and compact.

Theorem 1. The Todd genus of a manifold equipped with a symplectic circle action with isolated fixed points is either equal to zero and then the action is non-Hamiltonian, or equal to one and then the action is Hamiltonian. Any symplectic circle action on a manifold with the positive Todd genus is Hamiltonian.

Proof. A symplectic circle action is Hamiltonian if and only if there is such a connected component of the fixed point set that its weights of the S^1 -representation in the normal bundle are positive. This connected component is unique and corresponds to the global minimum of the moment map (see details in [1]). Let us apply the fixed point formula for Hirzebruch genus χ_y corresponding to the power series $\frac{x(1+ye^{-x(1+y)})}{1-e^{-x(1+y)}}$. Let M_s be connected components of the fixed point set, d_s be the number of negative weights of the S^1 -representation in the normal bundle of M_s , then (see [2])

$$\chi_y(M) = \sum_s (-y)^{d_s} \chi_y(M_s), \quad (1)$$

here the sum is taken over all connected components of the fixed point set. Since $\chi_y(pt) = 1$, the constant term of χ_y of a manifold equipped with a Hamiltonian circle action with isolated fixed points is equal to one. Vice versa, if the constant term of $\chi_y(M)$ is not equal to zero, then any symplectic circle action on M is Hamiltonian. The statement of the theorem follows from $Td(M) = \chi_y(M)|_{y=0}$.

The result of Theorem 1 is a particular case of the restrictions imposed on manifolds with a circle action by the Conner–Floyd equations. Let us fix

*The author was supported by Seggie Brown fellowship.

a multiplicative genus $h : \Omega_*^U \rightarrow Q$. Let $g_h(u) = u + \sum \frac{h([CP(n)])}{n+1} u^{n+1}$. For an almost complex manifold M^{2n} , equipped with a quasi-complex circle action with isolated fixed points, the sum over all the fixed points of the products $\prod_{k=1}^n (g_h^{-1}(j_k g(u)))^{-1} (j_k, k = 1, \dots, n, \text{ are weights of the } S^1\text{-representation in a neighborhood of the fixed point considered})$ contains no negative powers of u and its constant term is equal to $h(M^{2n})$ (see [3, 4]).

Corollary 1[6]. A semi-free symplectic circle action with isolated fixed points is Hamiltonian if and only if the fixed point set is non-empty.

Proof. Let us write the Conner-Floyd equations for the Todd genus in this case. Because $Td([CP(m)]) = 1$ for every m , we obtain $g_{Td}(u) = \sum u^{m+1}/(m+1) = \ln(1-u)$, $g_{Td}^{-1}(j g_{Td}(u)) = 1 - (1-u)^j$. All the weights of the S^1 -representation are equal to ± 1 . Summing up over all the fixed points we obtain that the function $f(u) = \sum_{k=1}^{2n} F_k \frac{(u-1)^k}{u^n}$, where F_k is the number of fixed points with exactly k negative weights, is a constant, say λ , which is equal to the Todd genus of the manifold. Solving the corresponding system of linear with respect to F_k equations, we find that $(F_0, F_1, \dots, F_n) = \lambda \left(\binom{n}{0}, \binom{n}{1}, \dots, \binom{n}{n} \right)$. Applying Theorem 1 we deduce the statement of the corollary.

Corollary 2[5]. A symplectic circle action on a 4-dimensional manifold is Hamiltonian if and only if the fixed point set is non-empty.

Proof. Let us show that the existence of a fixed point implies the existence of a fixed point whose weights of the S^1 -representation are of the same sign. Otherwise there is no connected component of codimension 2 (such component has only one weight). Thus, all fixed points are isolated and have exactly one negative and one positive weights. The contradiction follows from:

Lemma 1. Every quasi-complex circle action with isolated fixed points on M^{2n} has fixed points with different parity of the number of negative weights.

Proof. Otherwise the coefficients of $1/u^n$ in the Conner-Floyd equations for the Todd genus of the manifold have the same sign.

Let us consider in detail the case of a Hamiltonian circle action. Denote the Poincare polynomial of M by $P_M(t)$.

Theorem 2. If a symplectic manifold M admits a Hamiltonian circle action with isolated fixed points, then $\chi_y(M) = P_M(\sqrt{-y})$.

Proof. Critical points of the moment map coincide with the fixed points of the circle action. The index of every critical point is twice the number of negative weights of the S^1 -representation. Thus, the statement of the theorem follows from (1).

Corollary 3[7]. The signature and the Betti numbers of a manifold M^{4n}

equipped with a symplectic circle action with isolated fixed points are related through the formula: $\sigma(M^{4n}) = \sum_{k=0}^n b_{4k} - \sum_{k=0}^{n-1} b_{4k+2}$.

Proof. The statement follows from Theorem 2 after substitution $y = 1$.

Corollary 4. Homotopy equivalent manifolds equipped with Hamiltonian circle actions with isolated fixed points have the same χ_y -genus.

Remark 1. Similar to [8] for a non-Hamiltonian symplectic circle action with isolated fixed points $\chi_y(M^{2n}) = \sum_{k=0}^n b_{2k}(-y)^k$, here b_k are Novikov numbers [9] corresponding to the generalized moment map. It is important to construct an example of such an action with a non-empty fixed point set.

References

- [1] M. Audin // The topology of torus action on symplectic manifolds // Progr. in Math. 93, Birkhauser, 1991.
- [2] F Hirzebruch, T. Berger, R. Jung // Manifolds and modular forms// Aspects of Mathematics, E20, Braunschweig, 1992.
- [3] S. P. Novikov // Adams operators and fixed points // Izv. Akad. Nauk SSSR Ser. Mat. 32 (1968), 1245–1263.
- [4] I. M. Krichever // Formal groups and the Atiyah–Hirzebruch formula // Izv. Akad. Nauk SSSR Ser. Mat. 38 (1974), 1289–1304.
- [5] D. McDuff // The moment map for circle actions on symplectic manifolds // J. Geom. Phys. 5 (1988), no. 2, 149–160.
- [6] S. Tolman, J. Weitsman // On semi-free symplectic circle actions with isolated fixed points // Topology 39 (2000), no. 2, 299–309.
- [7] J. D. S. Jones, J. H. Rawnsley // Hamiltonian circle actions on symplectic manifolds and the signature // J. Geom. Phys. 23 (1997), no. 3–4, 301–307.
- [8] M. Farber // Signature in terms of Novikov numbers // Russian Math. Surveys 54 (1999), no. 1, 209–212
- [9] S. P. Novikov // The Hamiltonian formalism and a multivalued analogue of Morse theory (Russian)// Uspekhi Mat. Nauk 37 (1982), no. 5(227), 3–49.

University of Edinburgh
feldman@maths.ed.ac.uk